


USIRG's Frequently Asked Questions (FAQs)



TOPIC: Gelatin/Jell-O from the United States

BACKGROUND

IDDSI framework describes “*edible gelatin*” as a “sticky or gummy texture” (page 23-24 of https://iddsi.org/IDDSI/media/images/Complete_IDDSI_Framework_Final_31July2019.pdf), but the typical prepared gelatin products used in the United States (such as Jell-O brand gelatin) are different.

This gelatin/Jell-O in the USA is not the same as the Dysphagia Training Jelly of Japan. The gelatin/Jell-O of the USA is also very different from the Japanese konjac jelly, which is extremely gummy, cannot be pulled apart, and presents a very high choking risk.

This FAQ will address the prepared gelatin/Jell-O that is offered in many healthcare institutions and used in home settings around the United States. It may be homemade or bought ready-to-serve.

QUESTION 1: What IDDSI Level is prepared gelatin/Jell-O?

ANSWER:

Per testing and due to the challenging and **variable nature** of gelatin/Jell-O preparations across the USA, it is best practice to have gelatin/Jell-O only on the Regular, IDDSI Level 7 and Thin Liquid, Level 0 menus, unless:

- Your facility’s IDDSI testing indicates otherwise. This is based on your products, methods of preparation, and recipes. For example, gelatin/Jell-O may test appropriately for those on an Easy to Chew, Level 7.
- Provided based on a case-by-case assessment by a speech pathologist (SLP).
- Provided based on a Person-Centered Care approach (see PCC statement below).

The [IDDSI Testing Methods](#) yielded variable and contradicting results across many gelatin products. The following characteristics make gelatin/Jell-O a unique and potentially challenging food:

- Using the Spoon Tilt Test, it slides off spoon in one firm chunk.
- Using the Fork or Finger Pressure Test the results were variable, as samples:
 - Squashed with a fork, but broke apart into small particles, which will not form a cohesive bolus to meeting IDDSI testing standards, or
 - Remained too firm to squash between fingers.
- Upon opening a pre-packaged product, a separate layer of thin liquid was present.
- Using IDDSI’s Transitional Food testing ([see page 10 of Testing Methods](#)), gelatin/Jell-O does not melt to a thin liquid when water is applied.

QUESTION 2: Can we thicken the gelatin/Jell-O with a thickener product in the process of preparing it?


ANSWER:

Thickening gelatin/Jell-O can be difficult and may take trial-and-error until a palatable and safe product is attained. Reach out to your facility’s suppliers for recipes to try with various thickener products. There may also be commercially pre-packaged gelatin-like products that test to be appropriate for specific IDDSI levels. Always test all prepared or purchased products with the appropriate IDDSI Testing Methods to confirm the IDDSI level.

PERSON-CENTERED CARE (PCC) STATEMENT

The IDDSI Framework cannot change, but the diet order can be individualized. Person-centered diet orders specify an individual’s needs within and outside the IDDSI Framework. The medical team works in collaboration with the person’s wishes, preferences, goals of care, and individual abilities. Risks and benefits may be addressed in the informed consent process. Clear orders, communication, and documentation are key to person-centered care.

USIRG's Frequently Asked Questions (FAQs)



References for Gelatin and Japanese Training Jelly:

- Cichero, J. A. Y. (2020). Evaluating chewing function: Expanding the dysphagia field using food oral processing and the IDDSI framework. *Journal of Texture Studies*, 51(1), 56–66. <https://doi.org/10.1111/jtxs.12462>
- Kyodo, R., Kudo, T., Horiuchi, A., Sakamoto, T., & Shimizu, T. (2020). Pureed diets containing a gelling agent to reduce the risk of aspiration in elderly patients with moderate to severe dysphagia: A randomized, crossover trial. *Medicine (United States)*, 99(31), E21165. <https://doi.org/10.1097/MD.00000000000021165>
- Maksimenco, A., Lyude, A., & Nishiumi, T. (2020). Texture-modified foods for the elderly and people with dysphagia: Insights from Japan on the current status of regulations and opportunities of the high pressure technology. *IOP Conference Series: Earth and Environmental Science*, 548(2). <https://doi.org/10.1088/1755-1315/548/2/022106>
- Nishinari, K., Ishihara, S., Hori, K., & Fang, Y. (2020). Tongue-palate squeezing of soft gels in food oral processing. *Trends in Food Science and Technology*, 99, 117–132. <https://doi.org/10.1016/j.tifs.2020.02.023>
- Steele, C. M., Alsanei, W. A., Ayanikalath, S., Barbon, C. E. A., Chen, J., Cichero, J. A. Y., Coutts, K., Dantas, R. O., Duivesteyn, J., Giosa, L., Hanson, B., Lam, P., Lecko, C., Leigh, C., Nagy, A., Namasivayam, A. M., Nascimento, W. v., Odendaal, I., Smith, C. H., & Wang, H. (2015). The influence of food texture and liquid consistency modification on swallowing physiology and function: A systematic review. *Dysphagia*, 30(1), 2–26. <https://doi.org/10.1007/s00455-014-9578-x>
- Umay, E., Eyigor, S., Karahan, A. Y., Keskin, D., Karaca, G., Unlu, Z., Tikiz, C., Vural, M., Aydeniz, B., Alemdaroglu, E., Bilir, E. E., Yaliman, A., Sen, E. I., Akaltun, M. S., Altindag, O., Keles, B. Y., Bilgisoy, M., Ozcete, Z. A., Demirhan, A., & Gundogdu, I. (2019). Which swallowing difficulty of food consistency is best predictor for oropharyngeal dysphagia risk in older person? *European Geriatric Medicine*, 10(4), 609–617. <https://doi.org/10.1007/s41999-019-00206-z>